

The Commentary **printed in red** was written AFTER this article was published.

Published in the Military Times

March 9th, 2023

<https://www.militarytimes.com/off-duty/military-culture/2023/03/09/pentagon-ufo-chief-says-alien-mothership-in-our-solar-system-possible/>

MilitaryTimes

News Pay & Benefits Flashpoints Education and Transition Pentagon & Congress Off Duty Veterans Military Honor Subscribe Now



Pentagon UFO chief says alien mothership in our solar system possible

By **Zamone Perez**

Thursday, Mar 9



(Canva via Getty)

There is a possibility that extraterrestrial motherships and smaller probes may be visiting planets in our solar system, the head of the Pentagon's unidentified aerial

Featured Video



The future of Army stockpiles and Marine Bases | Defense News Weekly full episode 03.18.23



Will these USAF planes fall apart?



Watch Afghanistan vets' emotional testimony about the Afghanistan withdrawal



Will helping Ukraine diminish U.S. munition stockpiles?

Constraints on UAP

Draft version March 7, 2023

Typeset using L^AT_EX default style in AASTeX631

PHYSICAL CONSTRAINTS ON UNIDENTIFIED AERIAL PHENOMENA

Abraham (Avi) Loeb¹ and Sean M. Kirkpatrick²

¹*Head of the Galileo Project, Astronomy Department, Harvard University
60 Garden Street, Cambridge, MA 02138, USA*

²*Director of All-domain Anomaly Resolution Office
1010 Defense Pentagon
Washington DC 20301, USA*

ABSTRACT

We derive physical constraints on interpretations of “highly maneuverable” Unidentified Aerial Phenomena (UAP) based on standard physics and known forms of matter and radiation.

The title of this draft, Physical Constraints is most appropriate. Note the candid admission that the authors’ interpretation is restricted to limited, current day establishment or status quo accepted knowledge. This context is important to keep in mind as we consider this ET subject.

The authors are to be commended for openly approaching the subject of space traveling exo-astronauts. The following HISTORY is essential to this discussion: Two cases are key, the 1958 Jan 22nd CBS TV Armstrong Circle Theater (disclosure) broadcast that was pre-empted and censored by the USAF, and the 1966 Condon Report that considered the weakest cases, was made up of non-believers and designed to fool the public. In the name of national security, the pro-silence groups attacked the truth using debunking, lies, confusion and much more to bury this vital subject. This strategy continues today.

The combined authorship of this Harvard and Pentagon draft is welcome and intriguing in light of this history. In the spirit of those pioneers who pursued truth of this subject going back to the 1940’s and later we salute this one small step in the direction of disclosure and ongoing discovery which we hope will stay focused to benefit humankind on planet Earth and beyond. Note: The pro-silence groups, continuing their attacks to discredit, demean and debunk focus on confusing and destroying the ET component of this vital subject. This includes personal and professional attacks on Dr. Loeb, collaborators and predecessors. Like in the 40’s, 50’s and beyond you will see more of these attacks from those invested in protecting the status quo.

In particular, we show that the friction of UAP with the surrounding air or water is expected to generate a bright optical fireball, ionization shell and tail - implying radio signatures. The fireball luminosity scales with inferred distance to the 5th power. Radar cross-section scales similarly to meteor head echoes as the square of the effective radius of the sphere surrounding the object, while the radar cross-section of the resulting ionization tail scales linearly with the radius of the ionization cylinder. The lack of all these signatures could imply inaccurate distance measurements (and hence derived velocity) for single site sensors without a range gate capability.

Constraints on UAP

The lack of these signatures could also imply that our current status quo technology standards are not suited to measure technologies being used by highly advanced space traveling neighbors.

1. INTRODUCTION

In 2005, the US Congress tasked NASA to find 90% of all Near Earth Objects (NEOs) that are larger than 140 meters (Loff 2014). The Congressional task resulted in the construction of the Pan-STARRS telescopes. On October 19, 2017, the Pan-STARRS sky survey flagged an unusual NEO, the interstellar object 'Oumuamua (see, Loeb (2022a) and references therein). Unlike Solar system asteroids or comets, 'Oumuamua appeared to have an extreme flat shape and was pushed away from the Sun without showing a cometary tail of gas and dust, raising the possibility that it was thin and artificial in origin. Three years later, Pan-STARRS discovered a definitely artificial object, namely NASA's rocket booster 2020 SO, which exhibited similar behavior with an extreme shape, a push by the Solar radiation pressure and no cometary tail because its thin walls were made of stainless steel (Talbert 2020).

On March 9, 2017, six months before 'Oumuamua's closest approach to Earth, a meter-size interstellar meteor (IM2) collided with Earth (Siraj & Loeb 2022a). Surprisingly, IM2 had an identical speed relative to the Sun at large distances and an identical heliocentric semimajor axis as 'Oumuamua had. But the inclination of IM2's orbital plane around the Sun was completely different from 'Oumuamua's, implying that the two objects are unrelated.

Nevertheless, the coincidences between some orbital parameters of 'Oumuamua and IM2 inspires us to consider the possibility that an artificial interstellar object could potentially be a parent craft that releases many small probes during its close passage to Earth, an operational construct not too dissimilar from NASA missions.

Logical thinking is noted here with the "parent craft" narrative. This has been proposed, discussed and also demeaned and debunked since the UFO's began filling our skies in the 1940's. Hopefully this open and logical discussion continues without going too far down the rabbit hole of denial and ridicule that side tracked this vital subject over the past 70+ years.

These "dandelion seeds" could be separated from the parent craft by the tidal gravitational force of the Sun or by a maneuvering capability. A small ejection speed far away could lead to a large deviation from the trajectory of the parent craft near the Sun. The changes would manifest both in arrival time and distance of closest approach to Earth. With proper design, these tiny probes would reach the Earth or other Solar system planets for exploration, as the parent craft passes by within a fraction of the Earth-Sun separation - just like 'Oumuamua did. Astronomers would not be able to notice the spray of mini-probes because they do not reflect enough sunlight for existing survey telescopes to notice them if they are on the 10 cm scale of CubeSats or smaller. At a distance d from the Sun and the telescope, objects that are a meter in diameter and reflect a fraction $a \approx 10\%$ of sunlight impinging on their surface would yield a flux of optical light of $\sim 0.2(d/1 \text{ AU})^{-2} \text{ nJy}$, well below the detection threshold of even the *James Webb*

The referenced "seeds" have also been a serious part of academic and professional level discussion over the years. From Scout Ships to Sensing Discs we have evidence of ET visitations. The April 7th, 1952 Life Magazine delivered an historic disclosure of the ETs which was aided by Air Intelligence but was then denied and debunked. Look Magazine 4-1-52 was another. Washington Post 7-21-02 was a review of the documented (yes, denied and debunked) ET's flying over Washington DC.

The gigantic missing piece in this discussion is the HUMAN ET component. While the

Constraints on UAP

authors chose not to include the Human ET element here at this time, let us remember **September 1986: Dr. Cyril Ponnampereuma**, Dir. U. of Maryland's Laboratory of Chemical Evolution made headlines that read, **ET may look like us** – "It is demanded by chemistry - with the same elements believed to exist throughout the universe, there is a strong likelihood that the genetic code can spell out proteins only in the way known on Earth." "When we do land on a planet ... don't be surprised if somebody walks up to shake your hand." The fundamentals of life exist and surround us in space and this like organic matter has been raining down on our planet, and ALL planets, by tons each day, since the beginning he shared.

Space Telescope. In contrast, the radar signatures of a meter class object would be detectable with our deep space radars and space fence, much like IM2 was, out to beyond geosynchronous orbit at an altitude above 36,000 km. Such objects could also become optically detectable as they get close to Earth, especially if they create a fireball as a result of their friction with air.

There is overwhelming historical evidence of the Fire Balls. For one starting point, go back to the mid-February 1949 conference at Los Alamos where witnesses such as Dr's. La Paz, Teller, Kaplan and top military brass discussed this.

Equipped with a large surface-to-mass ratio of a parachute, technological "dandelion seeds" could slow down in the Earth's atmosphere to avoid burnup and then pursue their objectives wherever they land. Current radar coverage of the majority of first-world countries gives detectability of this High-Area-To-Mass (HAMR) objects down to a few centimeters depending on material, making detectability possible (Frueh et al. 2017).

Within a close range to a star, extraterrestrial technological probes could use starlight to charge their batteries and liquid water as their fuel. This would explain why they would target the habitable region around stars, where liquid water may exist on the surface of rocky planets with an atmosphere, like the Earth. Habitable planets would be particularly appealing to trans-medium probes, capable of moving between space, air and water. From a large distance, Venus, Earth or Mars would be equally attractive for probes. But upon closer inspection, Earth would show spectral signatures of liquid water (through reflection of blue light) and vegetation (through its red edge) that might attract selective attention (Seager et al. 2005).

What would be the overarching purpose of the journey? In analogy with actual dandelion seeds, the probes could propagate the blueprint of their senders. As with biological seeds, the raw materials on the planet's surface could also be used by them as nutrients for self-replication or simply scientific exploration. It is important to note, that given the time scales associated with the propulsion scheme discussed here, it is unreasonable to assert that the intention of any such probe launched in the far distant past, has anything to do with the human species. More likely, and similar to NASA's missions – the goal would be scientific and exploratory in nature.

"UNREASONABLE" "to have anything to do with the human species." Of course, just like the pro-silence groups over the past 70 years, there is tremendous pressure to shut the door on any and all HUMAN ET connection.

Keep the thinking tied into the traffic circle of technology as we know it today and hope that the majority are pre-occupied enough to forget, or not consider, that we are dealing with highly advanced space traveling civilizations. Advanced travelers who understand and use natural gravitational forces (as opposed to explosive or sling-shot motivation) to both propel and protect their voyages through space at speeds and maneuvers beyond the comprehension of some earthlings whose careers and livelihoods are locked into the status quo.

Constraints on UAP

Based on the detection rate of interstellar objects, [Siraj & Loeb \(2022b\)](#) estimated that for every interstellar NEO there are a thousand Solar system NEOs of the same size. Searching for interstellar meteorites among the many more meteorites from the Solar system without information about impact velocity, is like searching for a needle in a haystack.

This is why the first interstellar meteor (IM1), confirmed by velocity measurement of the US Space Command ([Siraj & Loeb 2022c](#)), is the target of a fully-funded ocean expedition by the Galileo Project ([Siraj et al. 2022](#); [Tillinghast-Raby et al. 2022](#)). Hopefully, by retrieving IM1's fragments within the coming year we will know whether its extraordinary material strength resulted from it being made out of an artificial alloy, like stainless steel or materials not yet developed by humans.

Are there any functioning extraterrestrial probes near Earth? We do not know. But the Galileo Project (2021) ([Loeb 2021](#)) intends to use the scientific method to explore this possibility, following the 2021 report about Unidentified Aerial Phenomena (UAP) from the Office of the Director of National Intelligence to the US Congress ([ODN 2021](#)). The state-of-the-art suite of instruments and computer algorithms of the Galileo Project will be able to study such data in the near future ([Loeb & Laukien 2023](#)).

"We do not know." The chronology of visitations attested to by pilots, radar technicians, CIA directors, generals, admirals, members of congress and other highly qualified people indicated very clearly, at least by their standards, WE DO KNOW.

The search for UAP, and the characterization of UAP, requires bounding the search plan with physics-based constraints on what we are searching for. This paper aims to constrain one aspect of the UAP hypothesis with parameters that govern the movement and interaction of a UAP with Earth's atmosphere to eliminate or bound observational uncertainties. Some data collected to date, while arising from multiple sensors, have uncertainties in one or more dimensions, leaving the exploitation of the data with a significant range of interpretations. This inevitably leaves open the debate on what some objects are, and whether or not they exhibited truly anomalous behavior. Specifically, if some observed UAP are of extraterrestrial origin, there are some practical limits on the interpretation of observed and measured data resulting from physics-based constraints.

2. THE EXTRATERRESTRIAL POSSIBILITY

The academic interest in UAP stems from their potential non-human technological origin. Extraterrestrial equipment could arrive in two forms: space trash, similar to the way our own interstellar probes (Voyager 1 & 2, Pioneer 10 & 11 and New Horizons) will appear in a billion years, or functional equipment such as autonomous devices equipped with Artificial Intelligence (AI). Electronic probes employing conventional chemical propulsion and refueling that we currently understand, would be a likely choice for travel within a planetary system. Some combination of conventional propulsion, ion propulsion, or lightsail propulsion would provide good choices for crossing the tens of thousands of light years that span the scale of the Milky Way galaxy. Such autonomous systems could be designed to survive even if the senders are not able to communicate with them, and deposit probes upon arrival to the target planetary systems. It is likely that any functional devices embedded in the Earth's atmosphere are not carrying biological entities because these would not survive the long journey through interstellar space and its harsh conditions, including bombardment by energetic cosmic-rays, X-rays and gamma-rays ([Hoang et al. 2017, 2018](#); [Hoang & Loeb 2020](#)).

"...not carrying biological entities because these would not survive the long journey." Remember the adage about repeating it enough times and then, that's all they think about. So, let's flip that coin and remind the readers that this report is based on locking the possibilities into the limitations of earth-based technologies as we know them today.

Interstellar gas and dust particles deposit a kinetic energy per unit mass that exceeds the output of chemical explosives at the speed of tens of km/s. However, technological devices with AI can be shielded

Constraints on UAP

to withstand the hazards of space, repair themselves mechanically, or even reproduce given the resources of a habitable planet like Earth. With Machine Learning capabilities, they can adapt to new circumstances and pursue the goals of their senders without any need for external guidance.

AI, ion propulsion, 3D printing, etc. are logical and somewhat creative applications of current day earth-based technologies. BUT this thinking locks us into what we know today and steers our truly creative thinking away from the very advanced standards already demonstrated by our space traveling ET neighbors. Such advanced technologies are being experimented with on earth, yes, with ET help (per Herman Oberth, Father of Modern Rocketry) but often such forward thinking and development encounters opposition.

As argued by John von Neumann in 1939, the number of such devices could increase exponentially with time if they self-replicate (Freitas 1980), a quality enabled by 3D printing and AI technologies. Physical artifacts might also carry messages, as envisioned by Ronald Bracewell in 1960Z (Bracewell 1960; Freitas & Valdes 1985) and currently used by NASA.

3. PROPULSION METHODS

In principle, the fastest devices could be launched by lightsails, pushed by powerful light beams up to the speed of light (Guillochon & Loeb 2015a). Natural processes, such as stellar explosions (Loeb 2023; Lingam & Loeb 2020) or gravitational slingshot near black hole pairs (Guillochon & Loeb 2015b; Loeb & Guillochon 2016), could launch objects to similar speeds. However, it would be difficult for relativistic payloads to slow down below the escape speed of Earth, $10^{-4.5}c$, without having around the same facilities that generated their high initial speeds.

A better-suited propulsion technique that was used in all interplanetary space missions from Earth is chemical rockets. Since rockets carry their fuel, they can navigate to a desired planet and slow down near it. Alternatively, it may be possible to use one of the above methods to travel to and through a planetary system, deploying interplanetary probes using conventional chemical propulsion. For a rocket of total mass, m , and exhaust speed of the ablated gas relative to the rocket, v_{exh} , momentum conservation implies: $m\dot{v} = -\dot{m}v_{exh}$, where an overdot, $(\dot{})$, denotes a partial time derivative. The

Tsiolkovsky solution to the rocket equation (Tsiolkovsky 2000), $(m_{initial}/m_{final}) = \exp\{(v_{final} - v_{initial})/v_{exh}\}$,

implies that for a reasonable fuel-to-payload mass ratio, the final speed v_{final} will only be an order of magnitude larger than the exhaust speed. For typical chemical propellants with v_{exh} of order a few km s^{-1} , this tyranny of the rocket equation explains why all human-made spacecraft reached a speed limit of tens of km s^{-1} or $\approx 10^{-4}c$. Interestingly, this speed is comparable to the escape speed from the

Earth's orbit around the Sun, $v_{esc} \approx 42\text{km s}^{-1}$, making it possible for humanity to launch interstellar

probes which take advantage of the motion of the Earth around the Sun at $v_{initial} \approx 30\text{ km s}^{-1}$. In contrast, chemical propulsion may not be sufficient for probes to escape from the habitable zone around dwarf stars, like the nearest star, Proxima Centuari (Loeb 2018; Lingam & Loeb 2018). In summary, chemical propulsion allows escape from the habitable zone of Sun-like stars and enables slowing down near a destination.

“...chemical propulsion may not be sufficient.” Absolutely true, and all the more reason to pull the shades away from the wealth of knowledge accumulated over the past 70 years, admit the history (yes, the mistakes too) and in the light of full disclosure, move forward.

Constraints on UAP

Devices which need to refuel would favor a habitable planet where liquid water or combustible organic fuel are available. The exhaust velocity of hydrogen/oxygen for rocket fuel is about 4.5 km s^{-1} (at a mass ratio of 16.4 with steam as exhaust) and so pure liquid hydrogen/oxygen is insufficient to slow down the characteristic free-fall speed of over 40 km s^{-1} from interstellar space to the habitable zone around the Sun unless the fuel mass is many orders of magnitude larger than the payload mass. Combustion of fuels with potentially higher heat capacity (such as hydrocarbons) would result in other chemical at the craft's exhaust, which would have distinct spectral signatures. This implies that chemical propulsion, while sufficient to escape the habitable zone using Earth's motion around the Sun, is insufficient to slow down from interstellar space to the planet's surface in the habitable zone without other assistance. Consequently, the mothership/probe scenario is more energetically viable. In addition, using water as the basis of the fuel would also require cold temperatures. Between hot exhaust (steam or other chemical byproducts) and cold storage (20K for hydrogen), this gives rise to additional signatures for characterization. Planets can be identified from a distance as they transit their star or through direct imaging (Winn 2023). Once an Earth-like planet is targeted, an interstellar device can plunge into its atmosphere. In principle, a multitude of tiny devices can be released from a mothership that passes near Earth. At $v_{final} \approx 10^{-4}c$, a probe would cross twice the distance of the Sun from the Milky-Way center within a time of $\approx 0.5 \text{ Gyr}$. The fraction of all Sun-like stars that host Earth-like planets in their habitable zone is in the range $\sim 3\text{--}100\%$ (Zink & Hansen 2019; Hsu et al. 2020; Bryson et al. 2021).

This implies that self-replicating probes could reach $\sim 10^{10}$ habitable planets around Sun-like stars in less than a billion years. Since most stars formed more than a billion years before the Sun (Madau & Dickinson 2014), it is possible that other technological civilizations predated ours by the amount of time needed for their devices to reach Earth.

“...self-replicating probe.” If we did not know of the technologies already in play by our space traveling neighbors, this would make some sense. So, if we are genuinely interested in pursuing real tangible answers, let's engage the wheels of something similar to a Manhattan Project to disclose, research, develop and apply these technologies.

Here we can use time as another constraint. In the propulsion scheme where an interstellar self-replicating autonomous system is travelling at $10^{-4}c$, the above analysis argues that $\sim 10^{10}$ habitable

planets around Sun-like stars could be reached within $\sim 0.5 \text{ Gyr}$. These self-replicating systems would necessarily be looking for water in order to generate fuel, and would necessarily have to take into account the relative motion of the planet in order to reach escape velocities after completion of the exploration mission. 1 Gyr ago Earth had water coverage and some simple algae plant life. In the extreme, detection of Earth 1 Gyr ago from a technological civilization near the center of the Milky Way 0.5 Gyr ago would be needed to decide to intentionally navigate here. In doing so, the navigators would need to plan for where the solar system would be located 1 Gyr in the future from their point of observation. Under such considerations, it becomes more likely that either: (i) such interstellar probes are the result of an unintended arrival to a planetary system; (ii) a technological civilization much closer to us than the center of the Milky Way; or (iii) an alternative propulsion scheme like the mothership/probe system is used. A detailed statistical analysis by Ezell & Loeb (Ezell & Loeb 2022) showed that the inferred abundance of probes is distinctly different in case of objects being targeted towards particular regions of the galaxy, specifically habitable zones containing planets. 'Oumuamua was detected at a distance of $\approx 0.2 \text{ AU}$ from Earth, and it passed through the habitable zone of our solar system. The estimated total number of 'Oumuamua-like objects would then fall by a factor of $\sim 2 \times 10^{10}$ in the case of targeted probes compared to

Not associated with the authors of this draft, the George Adamski Foundation & Space Science Educ. Proj. **commentary is noted in red**. Commentators: spacevoyagerllc @ gmail.com Cross reference to actual PDF from Harvard for correct formulas because PDF conversion to this document may have skewed the formulas.

Constraints on UAP

probes on random trajectories. The interstellar meteor IM1 had an estimated diameter of ~ 0.45 m and velocity of 60 km s^{-1} , but it was detectable when it burned up within the atmosphere of the Earth (Siraj & Loeb 2022a). The estimated detection rate for meter-size interstellar meteors is at least $\sim 0.1 \text{ yr}^{-1}$ (Siraj & Loeb 2022a), resulting in a local density estimate of $\sim 10^6 \text{ AU}^{-3} = 10^{22} \text{ pc}^{-3}$. This implies 8×10^{34} IM1-like objects bound by the thin disk of the Milky Way. However, if objects with the properties of IM1 were targeted towards habitable zones containing planets, the required number of such objects is merely $\sim 4 \times 10^{24}$. IM2 had a similar inferred number density to IM1 and a velocity of 40 km s^{-1} relative to the Local Standard of Rest (Siraj & Loeb 2022a). This implies $\approx 3 \times 10^{34}$ IM2-like objects, with a reduction to 1.5×10^{24} if such objects were targeted towards habitable zones.

The actual abundance of interstellar objects can be calibrated through future surveys such as the *Legacy Survey of Space and Time* (LSST) on the *Vera C. Rubin Observatory* in Chile. Parallax data from the James Webb Space Telescope may identify the nature and 3D trajectory of more 'Oumuamua-like or smaller interstellar objects crossing through or trapped within the solar system.

Below we show that any supersonic motion by such devices through the Earth's atmosphere would inevitably be accompanied by bright optical emission and detectable characterization signatures.

4. OPTICAL EMISSION

An object made of known matter with a frontal cross-sectional area A , moving at a supersonic speed, v , must create a bow shock in the Earth's atmosphere and dissipate a mechanical power

$$P \approx 12 A v^3 = 1.5 \text{ TW} (A/10 \text{ m}^2) (\rho_a/0.3 \text{ kg m}^{-3}) (v/10 \text{ km s}^{-1})^3, \quad (1)$$

where ρ_a is the ambient air density which depends on elevation, normalized here by a representative value at an altitude of 10 km.

Data on meteors shows that the fraction of the kinetic power which is radiated away in the optical band is $\approx 10\%$ [see equation (1) and figure 2 in Brown et al. (Brown et al. 2002)], implying an optical luminosity,

$$L_{\text{opt}} \approx 150 \text{ GW} (A/10 \text{ m}^2) (\rho_a/0.3 \text{ kg m}^{-3}) (v/10 \text{ km s}^{-1})^3. \quad (2)$$

For a path length l , this luminosity will persist over a period of time, $\sim 1 \text{ s} \times (l/10 \text{ km})/(v/10 \text{ km s}^{-1})$. Since $L_{\text{opt}} \propto A v^3$, the fireball luminosity scales with inferred distance to the 5-th power because A scales as distance squared and v scales as distance.

Encase a dual polarity multi-hulled ship in a protective self-generated gravitational force field and see what it appears to look like traveling through earth's atmosphere and beyond.

5. OTHER OBSERVABLE SIGNATURES

In addition to the thermal, shock, and associated optical signatures of a high velocity, highly maneuvering object, there is also an ionization and associated radio frequency signature from such an object moving through the atmosphere.

Studies into supersonic and hypersonic vehicles provide a good basis for comparison. While the ionization density depends on the altitude, shape, material and velocity of the object in motion, some limits can be derived on when a signature would be detected, implying a limit to the object's motion prior to the fireball threshold. In particular, ionization at high velocities leads to an increase in radar reflectivity along

Constraints on UAP

the ionization edge of the object and along the ionization trail. Both give rise to enhanced radio-frequency (RF) reflection for frequencies below the cutoff frequency.

Surzhikov calculates the ionization of air in high supersonic and low hypersonic regimes around blunt airframes (Surzhikov 2018). Figure 2 and table 1 in Surzhikov (2018) demonstrate the lowest end giving rise to a critical electron density of 10^{10} cm^{-3} for detection frequencies between 1-10 GHz. Variations of velocity, altitude and shape give rise to electron densities above this threshold, rendering the object detectable for typical radars in the L, S, C, and X radio bands.

Dhakal et al. conducted an excellent assessment of constraints on dark matter using radar meteor detectors (Dhakal et al. 2022). While not directly applicable to the current span of UAP sightings, the calculations do address smaller, faster objects at higher altitudes (70-130 km). These objects also exhibit an ionization trail which can be used for detection and measurement using similar radars. This regime is relevant to the mothership/probe scenario discussed above, indicating a detectable signature prior to the onset of a fireball as shown in Figures (1) and (11) of Dhakal et al. (Dhakal et al. 2022).

6. CONCLUSION

To put this vital discussion in what we consider a useful context, we urge an open recognition and understanding of the historical timeline into which this matter unfolds. There were, and still are, serious reasons why UN Sect. Gen. U Thant considered UFOs the most important problem facing the UN next to Vietnam. And why Vice Admiral Hillenkoetter, former CIA Director, would state that the UFO's are devices operating under intelligent control and that the truth is being kept from the public. This list of courageous pioneers goes on. And as the time goes on, we hope the authors of this report, their collaborators and successors earn their rightful place with these brave predecessors.

-1953 In a January 26th Department of Defense, Air Force Press Desk letter to Holt & Co, book publisher for Maj. Keyhoe, Ret. Director NICAP is stated: "We in the Air Force recognize Major Keyhoe as a responsible accurate reporter." "All the sightings and other information he listed have been cleared and made available to Major Kehoe from Air Technical Intelligence records, at his request." Read this **essential historical milestone**: Major Keyhoe's, [Flying Saucers from Outer Space](#) published by Holt & Co. 1953. His later, data-based books are built on this first book.

The considerations in this paper imply a useful limit on observations of UAP which bound the hypothetical explanations and can support limitations on interpretations of data. For example, one of the most common sets of data within the military holdings comes from FLIR (forward looking infrared) pods. These sensors provide an accurate resolved image of relative thermal measurements across the scene. Typical UAP sightings are too far away to get a highly resolved image of the object and determination of the object's motion is limited by the lack of range data. The range is usually estimated using the flight dynamics of the platform and some fixed points in the scene - if either are available. The error in estimating the range gives rise to a significant variation in the calculated velocity and is subject to human bias and error.

Claims of objects exceeding the transonic to supersonic range should be evaluated against the above known physics of ionization, radar reflectivity, temperature, sonic booms, and fireballs (Loeb 2022b). All of which can more effectively and accurately bound the velocity, and hence drive the range calculation. This will, in turn, when matched with the specifics of the sensor, allow for better estimates of the size,

Constraints on UAP

shape, and mass of the object in question.

CONGRESSIONAL Inquiry: Remember what 5 term U.S. Senator Barry Goldwater who completed his military career as a United States Air Force Major General was told when he requested ET evidence... **Request rejected "It's ABOVE TOP SECRET."**

Senator Goldwater was one of many whose ranks included other members of congress, CIA Director Vice Admiral Hillenkoetter, other admirals, generals, pilots, law enforcement, and many others who advocated disclosure and truth and faced the government's defiance and vicious obstruction.

ACKNOWLEDGEMENTS. This work was supported in part by Galileo Project at Harvard University and conducted in partnership with the Department of Defense, All-domain, Anomaly Resolution Office. We thank Richard Cloete for assistance and comments on the manuscript.

REFERENCES

- 2021, ODNI UAP Report], USG.
<https://www.dni.gov/files/ODNI/documents/assessments/Preliminary-Assessment-UAP-20210625.pdf>
Bracewell, R. N. 1960, Nature, 186, 670,
doi: 10.1038/186670a0
Brown, P., Spalding, R. E., ReVelle, D. O., Tagliaferri, E., & Worden, S. P. 2002, Nature, 420, 294,
doi: 10.1038/nature01238
Bryson, S., Kunimoto, M., Koppurapu, R. K., et al. 2021, AJ, 161, 36, doi: 10.3847/1538-3881/abc418
Dhakal, P., Prohira, S., Cappiello, C. V., et al. 2022, arXiv e-prints, arXiv:2209.07690,
doi: 10.48550/arXiv.2209.07690
Ezell, C., & Loeb, A. 2022, arXiv e-prints, arXiv:2209.11262, doi: 10.48550/arXiv.2209.11262
Freitas, R. A., J. 1980, Journal of the British Interplanetary Society, 33, 251
Freitas, R. A., J., & Valdes, F. 1985, Acta Astronautica, 12, 1027, doi: 10.1016/0094-5765(85)90031-1

DRAFT, UNDER REVIEW